



PAUL®
Laboratorio Veterinario

**VADEMECUM
PETS LINE**

✉ @LaboratorioPaul • Ⓛ Laboratorio Paul • ☎ 11 6304 8849
laboratoriopaul.com.ar • ventas@laboratoriopaul.com.ar

PAUL®
Laboratorio Veterinario

120
AÑOS
CUMPLIENDO!

PAUL®

Laboratorio Veterinario

PARTNERED IN
EVERY WAY



Committed to animal wellbeing since 1902.

INDEX

Ophtalmological Line

5. Colirio Paul

Antibiotic, Anti-inflammatory.

5. Diclovet

Anti-inflammatory.

6. Hyalmed

Treatment of erosive irritant conjunctivitis, ulcers without infectious involvement and dry eye.

6. Tobravet

Antibiotic.

7. Tobravet D

Anti-inflammatory, Intact.

7. Tacrolimus

Potent and Immediate Immunomodulatory Action.

Medical Line

10. Primevit 15

Vitamin Syrup

10. Paulhepat

Hepatoprotective Syrup

11. Coagulante Paul

Injectable. Hemostatic.

11. Lidocaína Paul

Injectable Solution.
Local Anesthetic

12. Paulvermín

Broad-Spectrum Internal
Antiparasitic

12. Flunixin Paul

Injectable Anti-inflammatory
Analgesic

Antibiotics

16. Enrofloxacina 5

Broad-Spectrum Antibacterial
Activity

16. Ceftiofur Paul

Broad-Spectrum Antibiotic Therapy

Otics

8. Paulderm

Otalgic - Anti-inflammatory
Antibiotic

8. Pauldex

Antibacterial - Antifungal - Anti-inflammatory

Biologics

13. Paulferón

Virostatic - Immunomodulator.

13. Paulvac antirrábica

Rabies vaccine

14. Paul 6

Six-in-One Vaccine

14. Parvo AT

Vaccine for the prevention of
parvovirus

15. Toxoidé Tetánico

Vaccine for prevention of
tetanus

15. PaulLepto

Vaccine for prevention of
leptospirosis

Massive Consumption

17. Silatox Plus Perro

External Antiparasitic Pipette

17. Silatox Plus Gato

External Antiparasitic Pipette

18. Espuma Seca

Biologically Active Shampoo

18. Silatox Dip

Cleansing Foam

19. Silatox S

Environmental Flea Killer

19. Silatox S Fogger

Environmental Flea Killer

20. Curabichera

Antimycosis, Healing, Bactericidal,
Fungicidal, Epithelializing, Repellent.

20. Shampoo Pulgues

External Antiparasitic

21. Shampoo Elastina

Shampoo for canine hygiene.

21. Shampoo Cachorro Dúo

Shampoo for puppy hygiene.

22. Shampoo Profesional

Shampoo for canine hygiene.

22. Dog Out

Repellent for Dogs and Cats

Dermatological

9. Paulderm

Antibiotic - Antifungal
Topical Anti-inflammatory

9. Pauloreja

For Ear Tip.
Antilarvic - Antiseptic
Repellent



OPHTHALMOLOGICAL LINE



Ophthalmological Line

COLIRIO PAUL



Formula:

- Neomycin sulfate: 0.886 g
- Betamethasone 21-sodium phosphate: 0.100 g
- Excipients q.s. to 100 mL

Presentation: 10 mL

Action:

- Antibiotic
- Anti-inflammatory

This combination provides a potent effect by combining Betamethasone 21-sodium phosphate, a powerful corticosteroid that controls inflammatory and exudative manifestations, with Neomycin sulfate, a broad-spectrum antibiotic.

- Free from cytotoxic preservatives.
- Isotonic formulation with physiological pH.

Indicated for the treatment of conjunctivitis, keratitis, keratoconjunctivitis, blepharitis, iritis, and iridocyclitis in dogs and cats.

Dosage and Administration:

Instill 1 to 3 drops in the conjunctival sac every 8 hours for 72 hours. The duration of treatment may be extended at the discretion of the attending veterinarian.

Contraindications:

- Do not use in patients with corneal ulcers.
- Do not use in patients with glaucoma.

DICLOVET



Formula:

Each 100 mL contains:

- Sodium Diclofenac: 0.100 g
- Sodium Hyaluronate: 0.200 g
- Excipients q.s.

Presentation:

Sterile ophthalmic drops / 5 mL

Contraindications:

Active epithelial keratitis caused by herpes simplex.

Dosage and Administration:

Unless otherwise directed by a veterinarian, apply 1 to 2 drops to the affected eyes 4 to 5 times a day, depending on the severity of the condition.

Action:

Non-steroidal anti-inflammatory for dogs and cats. Common indications include conjunctivitis, keratitis, keratoconjunctivitis, corneal ulcers, traumatic inflammation, and ocular pain.

Non-steroidal anti-inflammatory drugs (NSAIDs) reduce protein release from uveal vessels mediated by prostaglandins, diminishing the presence of antibodies that contribute to immunological and fibrin reactions causing synechiae. They also reduce inflammation-mediated miosis. Recommended when the use of corticosteroids is not advisable.

Ophthalmological Line

HYALMED



Formula:

Each 100 ml contains:

- Sodium Hyaluronate 0.200 g
- Excipients q.s.

Presentation: 10 ml

Action:

Physiologically balanced artificial tears. Acts as a humectant, lubricant, and epithelium-regenerating agent for the cornea and conjunctiva. It forms a protective film due to its viscosity and adhesion to the ocular surface.

Indicated for the treatment of irritative and erosive conjunctivitis, non-infectious ulcers, and dry eye.

Dosage and Administration:

Instill 1 to 2 drops in the affected eye two or three times a day, or as directed by the attending Veterinarian, until symptoms subside.

TOBRAVET



Formula:

Each 100 ml contains:

- Tobramycin base 0.300 g
- Sodium Hyaluronate 0.200 g
- Excipients q.s.

Presentation: 10 ml

Action:

Antibiotic and healing agent for the topical treatment of bacterial eye infections, corneal ulcers, corneal erosion, conjunctivitis, keratoconjunctivitis, keratitis, and blepharitis caused by susceptible bacteria.

Dosage and Administration:

Instill 1 or 2 drops into the conjunctival sac of the affected eye at 4- to 6-hour intervals until symptoms resolve or as directed by the attending veterinarian.

Contraindications:

Hypersensitivity to aminoglycosides administered topically may occur in some patients. If any hypersensitivity reaction occurs, discontinue use immediately.

Ophthalmological Line

TOBRAVET D



Formula:

Each 100 ml contains:

- Tobramycin base 0.300 g
- Dexamethasone base 0.100 g
- Sodium Hyaluronate 0.200 g
- Excipients q.s.p.

Presentation: 5 ml

Action:

Antibacterial, anti-inflammatory, and healing agent. Indicated for the topical treatment of bacterial eye infections causing conditions such as conjunctivitis, keratoconjunctivitis, keratitis, and blepharitis.

The addition of Dexamethasone provides potent anti-inflammatory and decongestant properties with excellent penetration and immediate effect. Meanwhile, Sodium Hyaluronate helps maintain corneal epithelial integrity and promotes wound healing.

Dosage and Directions for Use:

Instill 1 or 2 drops into the conjunctival sac of the affected eye at 4- to 6-hour intervals during the first 24 to 48 hours. Gradually reduce the frequency of application until the infection resolves, as per the attending veterinarian's instructions.

Contraindications:

- Do not use in viral infections of the cornea and conjunctiva, fungal infections, or corneal ulcers.
- Prolonged treatment may cause superinfection by non-susceptible microorganisms, including fungi.

TACROLIMUS



Formula:

- Tacrolimus monohydrate 3.07 mg
- Excipients q.s.p. 10 ml

Presentation: 10 ml

Indication:

Powerful and immediate immunomodulatory action.

Indicated for the treatment of immune-mediated ocular conditions with corneal and/or conjunctival lymphocyte and plasma cell infiltration, such as:

- Keratoconjunctivitis sicca (dry eye syndrome)
- Chronic superficial keratitis
- Nodular and necrotizing episcleritis in dogs and cats.

Acts as a lacrimal stimulant by reducing the immune response in autoimmune inflammation of the lacrimal gland, restoring natural tear production.

Dosage and Directions for Use:

Ready-to-use ophthalmic solution. Apply 1 drop every 12 hours (2 drops/day) for the period prescribed by the veterinarian, typically 15 to 30 days.

Its immediate action and excellent penetration into the cornea and conjunctiva ensure a potent immunomodulatory and anti-inflammatory response, providing relief and improving quality of life. Treatment duration may be adjusted at the discretion of the attending veterinarian.

Contraindications:

- Hypersensitivity to the active ingredient.
- Active ocular infections.

PAULDERM



Formula:

- Chloramphenicol base 1.45 g
- Triamcinolone acetonide 0.10 g
- Lidocaine hydrochloride 1.50 g
- Boric acid 5.00 g
- Excipients q.s.p. 100 ml

Presentation: 50 ml

Action:

Antibiotic, anti-inflammatory, and antifungal. The formulation effectively dissolves cerumen and secretions, ensuring proper delivery of active ingredients to the affected area.

Anti-inflammatory: Potent action promotes the atrophy of sebaceous glands, often associated with chronic otitis.

Antifungal: Boric acid eliminates *Malassezia pachydermatis*.

Alcohol-based solution: Enhances desiccation of the ear canal, creating unfavorable conditions for microorganism replication.

Indications:

- Treatment of chronic otitis
- Dermatitis of the external auditory canal
- Otitis externa
- Chronic suppurative otitis media
- Analgesic and antiseptic purposes

Dosage and Directions for Use:

Apply the equivalent of 1 or 2 pumps into the affected ear, as directed by the veterinarian.

PAULDEX



Formula:

- Chloramphenicol 0.50 g
- Benzocaine 1.00 g
- Dexamethasone 0.02 g
- Chlorobutanol 0.40 g
- Propylene glycol q.s.p. 100 ml

Presentation: Dropper bottle, 20 ml

Dosage and Directions for Use:

Apply 2 to 3 drops into the external ear canal 2 to 3 times per day, massaging the base of the ear afterward to ensure proper penetration and distribution.

For chronic cases, Paulderm Ótico is recommended.

Action:

Antibacterial, antifungal, and anti-inflammatory.

Chloramphenicol: Broad-spectrum antibiotic effective against gram-positive and gram-negative bacteria.

Dexamethasone: A corticosteroid with potent anti-inflammatory and antiallergic effects, reducing exudation, erythema, and itching.

Chlorobutanol and Propylene Glycol: Act as bacteriostatic, bactericidal, and fungicidal agents. Propylene glycol also reduces edema due to its hygroscopic action.

Benzocaine: Provides immediate local anesthetic relief.

Indications:

Acute otitis externa and media

Otic furunculosis

Dermatological

PAULDERM



Formula (per 100 ml):

- Gentamicin sulfate 0.10 g
- Triamcinolone acetonide 0.10 g
- Griseofulvin 1.50 g

Presentation: 50 ml

Action:

Antibiotic, anti-inflammatory, and antifungal.

This dermal gel combines:
Gentamicin: A broad-spectrum bactericidal antibiotic effective against bacterial infections.
Triamcinolone: A corticosteroid with potent anti-inflammatory, antiallergic, and vasoconstrictive properties.
Griseofulvin: A powerful fungistatic agent effective against dermatophytes such as *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton*.



Indications:

Treatment of skin and auricular pavilion infections, including:

- Pyoderma of folds
- Interdigital intertrigo
- Impetigo
- Bacterial folliculitis
- Dermatophytosis
- Hypersensitivity lesions
- Lick granuloma

Dosage and Directions for Use:

Apply topically 2 to 3 times daily to the affected area, rubbing gently to promote product penetration.

The duration of treatment is based on the remission of symptoms and at the discretion of the attending veterinarian.

PAULOREJA



Formula (each 100 ml contains):

- Permethrin 2.00 g
- Benzalkonium chloride 0.10 g
- Excipients q.s. 100 g

Presentation: 60 g

Action:

- Larvicidal, antiseptic, and fly repellent.
- Effective against myiasis (fly larvae infestation) in ear tips, tail tips, wounds, and ulcers susceptible to fly strikes.

Indications and Directions for Use:

For use in dogs only.

As a larvicide: Apply directly to wounds with myiasis or wounds on ear tips.

As a preventative: Apply to wounds and ulcers prone to fly infestation, especially during fly season.

Reapply daily.

Precautions:

- Do not apply to dogs younger than 4 weeks of age.
- Avoid use on weak or ill animals.
- Do not apply to the eyes, mouth, muzzle, or genitals.

PRIMEVIT 15



Formula:

- Folic acid
- Vitamin B1 (Thiamine pyrophosphate)
- Vitamin B2 (Riboflavin)
- Vitamin B6 (Pyridoxine hydrochloride)
- Vitamin B12 (Cyanocobalamin)
- Nicotinamide
- Vitamin B15 (Pangamic acid)
- Glucuronic acid
- Calcium pantothenate
- Acetyl DL-methionine
- Choline chloride
- Inositol

Presentation: 100ml

Action:

- Vitamins
- Lipotropic
- Reconstitutive
- Detoxifying
- Hepatonormalizing
- General tonic
- Appetite stimulant
- Central nervous system stimulant
- Antianemic

Dosage and Administration:

Administer 5 ml per 15 kg body weight 2 to 3 times daily via oral syringe

Indications:

- Complex vitamin B deficiencies
- Conditions of anorexia and malnutrition
- Weakness due to infectious or parasitic diseases
- Convalescence
- Adjunct to antibiotic therapy
- Metabolic imbalances with liver dysfunction (complementary therapy to Paulhepat)
- Neuritis and polyneuritis
- Gastrointestinal disorders

Paulhepat



Formula:

- Choline bitartrate
- Acetyl DL-methionine
- Inositol
- Cyanocobalamin (Vitamin B12)

Presentation: 100 ml - Syrup

Action:

- Hepatoprotective
- Treatment for hepatic dysfunction
- Infectious hepatitis
- Toxic hepatitis
- Medicamentous hepatitis
- Hepatic cirrhosis
- Hepatic origin dermatitis
- Hepatic origin urticaria
- Dyskinesia in dogs and cats

Dosage and administration:

- Administer orally 1 ml/kg body weight twice daily.
- The duration of treatment depends on the veterinarian's assessment.
- PAULHEPAT is a complementary therapy for treating liver diseases of various origins. The dosage should be tailored to meet the specific needs of each case, as determined by the veterinarian.

COAGULANTE PAUL



Formula:

- Oxalic acid 1.200 g
- Succinic acid 0.300 g
- Malic acid 0.300 g
- Sodium menadione bisulfite 0.300 g
- Lidocaine hydrochloride 0.300 g

Presentation: 10 ml and 50 ml

Action:

- Hemostatic.
- Indicated as both a prophylactic and curative treatment for surgical interventions.
- Useful for lateral or bilateral epistaxis, hemoptysis, hematemesis, metorrhagia, post-partum hemorrhage, surgical and traumatic bleeding, and hemorrhagic enteritis.

Dosage and Administration:

- Based on the severity of the case and the veterinarian's recommendation, administer 3-6 mg of sodium menadione bisulfite per 10 kg every 24 hours.
- Approximate dosing for each species:
 - Equines and bovines: 10 ml
 - Caprines, ovines, and swine: 2-5 ml
 - Felines and canines: 1-2 ml

Route of administration:

Intramuscular

- Administer every 12 to 24 hours for 3 to 5 days, or until coagulation time returns to normal.
- If symptoms persist, consult with the veterinarian to review the treatment plan.
- For external wounds: Apply Coagulante Paul on a gauze or cotton dressing.

LIDOCÁINA PAUL



Para pequeños y grandes animales



Formula:

Lidocaine hydrochloride 2.00 g

Presentation: 100ml

Action:

Local anesthetic, used for anesthesia and analgesia through infiltration, nerve conduction blockage, regional anesthesia, epidural anesthesia, and topical anesthesia.

Dosage and Administration:

The dosage should be determined by the attending veterinarian. For epidural injection, the general usage is:

Canines and Felines:

- Epidural anesthesia/analgesia: 1 ml (20 mg) per 3.5 to 4.5 kg of body weight (administered slowly)
- Epidural anesthesia: 5 to 10 ml (100 to 200 mg)

Equines and Bovines:

- Epidural anesthesia: 5 to 10 ml (100 to 200 mg)
- Nerve blockade: For example, 10 to 20 ml for maxillary nerve block; smaller amounts for other nerves.

FLUNIXIN PAUL



Fórmula:

- Flunixin 5.000 g
- Meglumine
- Excipients

Presentación: 20-50-250ml

Action:

- Anti-inflammatory
- Analgesic
- Antipyretic

Used for the treatment of:

- Colic
- Endotoxemia
- Musculoskeletal pain
- Arthritis
- Spinal disc injuries

Dosage and Administration:

Route: Intramuscular

Canines:

- For the treatment and control of musculoskeletal inflammatory processes.
- Dose: 0.5–2 mg/kg body weight, once a day or divided into two applications every 12 hours, for up to 3 consecutive days.

Bovines:

- For the treatment and control of musculoskeletal inflammatory processes, fever associated with respiratory diseases, and endotoxemia.
- Dose: 1–2 mg/kg body weight, once a day or divided into two applications every 12 hours, for up to 3 consecutive days.

Equines:

- For the treatment and control of musculoskeletal inflammatory processes and pain associated with colic.
- Dose: 1 mg/kg body weight, once a day for up to 5 consecutive days.
- For colic: Administer intravenously. If symptoms persist, the dose can be repeated.

Biologics

PAULFERÓN



Formula:

- Interferon alpha 2b: 30 IU

Presentation: 30ml

Action:

- Virostatic
- Immunomodulator

Paulferón contains recombinant human interferon alpha 2b, a protein produced using recombinant DNA techniques in *Escherichia coli* bacteria.

Indicated for the treatment of:

- Ocular disease attributed to herpesvirus
- Gingivitis, stomatitis, chronic pharyngitis
- Feline leukemia virus infection
- Supportive treatment for animals with retrovirus-associated tumors
- Neoplasms and paraneoplastic syndromes

Dosage and Administration:

- Herpesvirus-associated ocular disease:
 - 0.2 IU/kg orally, every 24 hours for 7 days
- Gingivitis, stomatitis, chronic pharyngitis:
 - 30 IU total. 1 ml orally every 12 hours for 7 days. Repeat the regimen after a one-week break
- Feline leukemia virus infection:
 - 30 IU total. 1 ml orally every 12 hours for 7 days. Repeat the regimen after a one-week break
- Supportive treatment for retrovirus-associated tumors:
 - 30 IU total. 1 ml orally every 12 hours for 7 days. Repeat the regimen after a one-week break

Contraindications:

- Patients with a history of allergy to interferon, glycine, or albumin
- Epileptic disorders or functional nervous system alterations
- Severe cardiovascular disease
- Severe renal or hepatic dysfunction

PAULVAC ANTIRRÁBICA



Formula:

- Inactivated rabies virus suspension with Betapropiolactone, Pasteur strain developed in BHK 21 clone 13 cells, adjuvanted with aluminum hydroxide.
- Contains at least 1 IU of potency per milliliter.

Dosage and administration:

- 1 ml subcutaneously, starting at 3 months of age.
- Repeat annually.

Presentation: 10 dosis

Action:

Prevention of Rabies in dogs and cats (Monodose).

Biologics

PAUL 6



Formula:

Lyophilized Vaccine Components:

- Canine Distemper
- Canine Adenovirus Type II
- Canine Parainfluenza
- Canine Parvovirus (live attenuated).

Diluent:

- Inactivated Canine Coronavirus suspension

Presentation: 12 doses

Action:

Prevents canine distemper, hepatitis, kennel cough, parainfluenza, parvovirosis, and canine coronavirus infections.

Dosage and Administration:

- Administer 1 ml intramuscularly or subcutaneously after reconstitution.
- Puppies aged 45 days or older: Administer 3 doses at intervals of 3 to 4 weeks.
- If vaccinated before 4 months of age, a single booster dose should be given at 4 months due to potential interference from maternal antibodies.
- Annual revaccination with a single dose is recommended.

Precaution:

- Only vaccinate dogs in good health.

PARVO AT



Formula:

Suspension of modified live canine parvovirus.

Presentation: 24 doses de 1 ml

Action:

Vaccine for the prevention of canine parvovirosis.

Dosage and administration:

- Vaccinate dogs in good health only.
- Administer 1 ml intramuscularly or subcutaneously.
- Puppies aged 45 days or older: Administer 3 doses at intervals of 3 to 4 weeks.
- Puppies vaccinated before 5 months of age should receive a single booster dose at 5 months to counteract possible interference from maternal antibodies.
- Annual revaccination with a single dose is recommended.

Biologics

TOXOIDE TETÁNICO



Formula:

Inactivated suspension of Clostridium tetani toxoids, with aluminum hydroxide as an adjuvant.

Presentation: 20 mL or 4 mL

Action:

Vaccine for the prevention of tetanus caused by Clostridium tetani toxins in equines, ovines, porcines, and canines.

Dosage and administration:

Equines and Bovines: 2 mL.

Ovines, Porcines, and

Canines: 1 mL.

Administer via subcutaneous or intramuscular injection using a sterile syringe and needle.

Administer 60 and 30 days before parturition in primovaccinated animals. For initial vaccination (primovaccination): Apply two doses with an interval of 20 to 25 days, starting at 3 months of age.

Annual revaccination: Apply a single dose 30 days before parturition.

Depending on the professional's judgment, revaccination is recommended 1 or 2 times per year.

PaulLepto



Formula:

Cultures of *Leptospira interrogans* serogroups: Canicola strain Moulton and Icterohaemorrhagiae strain CF-1, inactivated, contains penicillin, streptomycin and amphotericin B as preservatives.

Presentation: 10 doses of 1ml

Action:

Vaccine for the prevention of canine leptospirosis.

Dosage and method of use:

1 ml. In animals that have been vaccinated for the first time, a booster is recommended 3 or 4 weeks after the first dose.


Then revaccinate annually or half-yearly according to the rate of risk of contagion and the criteria of the acting professional. It can be used as a vaccine diluent.

Contraindications:

Do not vaccinate animals with viral, bacterial or parasitic diseases.

Antibiotics

ENROFLOXACINA 5 PAUL



Formula:

Enrofloxacin 5 g.

Presentation: 100 mL vial.

Dosage and Administration:

Canines:

- Dosage: 0.5 to 1 mL per 10 kg of body weight.
- Route: Intramuscular injection.
- Administer every 12 hours.
- Continue treatment for 2–4 days after clinical signs subside, with a maximum duration of 10 consecutive days.
- If no improvement is observed within the first 5 days, re-evaluate the diagnosis and consider an alternative therapy.

Cattle, Swine, Sheep, and Goats:

- Dosage: 5 to 10 mL per 100 kg of body weight.
- Route: Subcutaneous injection.
- Administer every 24 hours.
- Duration of treatment: 3 consecutive days. May extend for 2 additional days in animals showing clinical improvement but incomplete recovery.
- Do not administer more than 20 mL per injection site.

Note:

- Local injection may cause a transient reaction, potentially resulting in tissue loss in edible areas at slaughter.

CEFTIOFUR PAUL



Formula:

- Ceftiofur 50 mg/mL
- Third-generation cephalosporins

Presentation:

100 mL – Ready-to-use injectable solution

Action:

- Broad-spectrum antibiotic therapy.
- Ceftiofur belongs to the third-generation cephalosporins, effective against Gram-positive and Gram-negative bacteria, including beta-lactamase-producing strains.



Dosage and Administration:

Cattle:

- Route: Intramuscular or subcutaneous injection.
 - Respiratory disease: 2 mL per 50 kg body weight. Repeat after two days.
 - Foot rot (Pietin): 1–2 mL per 50 kg body weight. Administer once daily for 3–5 consecutive days.

Swine:

- Route: Intramuscular injection.
 - Respiratory disease: 0.6–1 mL per 10 kg body weight. Administer once daily for 3 consecutive days.

Massive Consumption

SILATOX PLUS dogs



Formula:

- Fipronil
- Methoprene
- Formulation agents q.s.p.

Presentation: Spot-on pipette

Action:

- External antiparasitic pipette (flea treatment) with high knockdown power and excellent residual efficacy.
- Also effective against eggs and larvae.
- Effective against ticks and lice.

Dosage and Administration:

Apply topically to dry skin with the animal standing, on the back between the shoulders.

Recommended usage: 1 application per month.

Dosage by weight:

- Dogs over 8 weeks of age up to 10 kg: Pipette of 0.67 mL.
- Dogs 10–20 kg: Pipette of 1.34 mL.
- Dogs 20–40 kg: Pipette of 2.68 mL.
- Dogs 40–60 kg: Pipette of 4.68 mL.



SILATOX PLUS cats



Formula:

- Fipronil
- Methoprene
- Formulation agents q.s.p.

Presentation: Spot-on pipette

Action:

- External antiparasitic pipette (flea treatment) with high knockdown power and excellent residual efficacy.
- Effective against eggs and larvae.
- Also works against ticks and lice.

Dosage and Administration:

• Apply topically to dry skin with the animal standing, on the back between the shoulders.

• For animals aged 8 weeks or older and weighing up to 5 kg: Pipette of 0.5 mL.

• For animals weighing from 5 kg to 8 kg: Pipette of 0.8 mL.

• Recommended usage: 1 application per month.



Massive Consumption

ESPUMA SECA



Formula:

Sodium Lauryl Sulfate: 7.70 g

Presentation: 250 ml

Action:

Biologically active shampoo, ideal for washing dogs and cats of all ages. Effectively cleans while restoring shine and softness, without the need to wet the animal. Does not irritate the eyes or cause adverse reactions.

Dosage and Directions for Use:

Shake the container well and apply directly to the animal's body. Rub gently by hand until the foam disappears. Dry thoroughly and brush the coat.

SILATOX DIP



Formula:

Permethrin 0.50 g

Metoprene 0.25 g

Piperonyl butoxide 1.50 g

Presentation: 250 ml

Action:

Cleaning foam and external antiparasitic for the control and treatment of flea infestations in dogs. Also controls lice and ticks.

Dosage and Mode of Use:

Shake the container and apply to the animal's body. Then rub with your hand until the foam disappears. It is recommended to do this against the fur to ensure better penetration. Avoid contact with eyes and mucous membranes.



Massive Consumption

SILATOX S



Formula:

Permethrin 0.50 g
Metoprene 0.15 g
Piperonyl butoxide 1.50 g

Presentation: 440ml

Action:

Environmental flea control. Silatox S spray is indicated for the control, elimination, and prevention of flea infestations in areas where animals reside. It has a dual effect, eliminating adult fleas and preventing the hatching of eggs, maturation of larvae, and pupae of fleas.

Dosage and Mode of Use:

The content of Silatox S covers approximately 100 m². Apply in areas frequented by cats and dogs, including carpets, bedding, and furniture. It is ready to use. After application, it is recommended to leave the area for at least 2 hours and ventilate.

Precautions:

Ventilate the area. Carefully read the label on the container.

SILATOX S FOGGER



Formula:

Permethrin 0.50 g
Metoprene 0.15 g
Piperonyl butoxide 1.50 g

Presentation:

250 ml Aerosol with total discharge valve

Indications:

Environmental insecticide. For the control and elimination of fleas in areas where animals reside. Due to its composition, it has a dual effect: it eliminates adult fleas and prevents the hatching of eggs, as well as the maturation of larvae and pupae of fleas.

Precautions:

Ventilate the area. It is recommended to carefully read the product label.

Dosage and Usage:

The content of Silatox S covers approximately 70 m². It should be applied in areas frequented by felines and canines. It is ready to use. To achieve effective results in the treated areas, use the product as indicated. Close exterior doors and windows, turn off ventilation systems, open interior doors and furniture. Remove blankets, bedding, and carpets to be treated. Remove animals, aquariums, food, and feeding utensils. Turn off any lit flames, except for pilot lights. Place the Silatox container on a flat surface on a piece of paper, press the valve until it locks, keeping the container away from your body, and place it in the designated area. Leave the treated area and stay away for 2 hours. After 2 hours, ventilate for 30 minutes by opening doors, windows, and ventilation systems.

Massive Consumption

CURABICHERA



Formula:

Silver sulfadiazine
Dichlorvos
Cypermethrin
Aluminum

Presentation: 440ml

Dosage and Usage:

Spray onto the wound and its perimeter, forming a highly adhesive layer.
As a curative: apply once a day for 3 or 4 days.
As a preventive: a single application.

Action:

Anti-mite, healing, bactericidal, fungicidal, epithelializing, repellent.

SHAMPOO PULGUEX



Formula:

- Tetramethrin
- Cyclopropyl carboxylate
- Tetrahydronaphthalimide
- Piperonyl butoxide
- Formulation agents c.s.p.

Presentation: 30 ml

Action:

External Antiparasitic – Flea and Tick Killer.
Specially formulated for canines with insecticidal power by contact, providing immediate effect through foamy bathing.

Massive Consumption

SHAMPOO ELASTINA



Formula:

Sodium Laureth Sulfate
Cocamide DEA (Cocoyl
Diethanolamide)

Action:

Promopet Shampoo with Elastin has an exclusive formula that effectively cleans, restoring shine and softness.

Presentation: 330ml

SHAMPOO CACHORRO DUO



Formula:

- Sodium Laureth Sulfate
- Cocamide DEA (Cocoyl Diethanolamide)
- Oleic Alcohol
- Formulation Agents c.s.p.

Presentation: 330ml

Action:

Shampoo Puppy Duo is specially formulated for the hygiene of puppies of all breeds. It cleans and detangles simultaneously. Due to its formula and balanced pH, it does not irritate the eyes and can be used frequently. Its rich and abundant foam acts immediately, leaving the coat soft and shiny.

Massive Consumption

SHAMPOO PROFESIONAL



Formula:

- Sodium Laureth Sulfate
- Cocamide DEA (Cocoyl Diethanolamide)
- Formulation Agents c.s.p.

Presentation: 1lt

Acción:

Shampoo specially formulated for the hygiene of dogs of all breeds. Due to its formula and balanced pH, it does not irritate the eyes and can be used frequently. Its rich and abundant foam acts immediately, leaving the coat soft and shiny.

DOG OUT



Formula:

- Aerosol: Methyl Nonyl Ketone
- Formulation Agents
- Liquid: Undecanal

Presentation liquid: 500 ml

Presentation spray: 440 cm³

Dosage and Mode of Use:

Wash the area to be treated to remove any feces. Dilute 100 cm³ in 1 liter of water, applying by thoroughly wetting the area.

For the spray, spray from a distance of 50 cm.

Apply daily until the animals' behavior changes.

Action:

Repellent solution for dogs and cats (behavior modifier), containing 2-Undecanal as the active ingredient, along with other inert formulation agents, to be applied in the animal's environment.